## **Scintillation Kinetics through DFT**

## Electronic Structure Analysis to Understand Scintillation Kinetics of CsI Scintillator Single Crystal



(a) A CsI single crystal developed in BARC.

(b) Local crystal structure of CsI:Tl, CsI:Tl, Eu and CsI:Tl, Sr.

(c) Calculated partial density of states (PDOS) of CsI:TI (Top) and defect complex and  $TI_{cs}^{+}+Eu_{cs}^{+}$  (middle) and  $TI_{cs}^{-}+Sr_{cs}^{+}$  (bottom) in CsI matrix.

Dr. Shashwati Sen is currently heading Crystal Technology S e c t i o n o f Technical Physics Division, BARC. Dr. Sen specializes on the growth of scintillator and semiconductor



semiconductor single crystals for their application in radiation detection. She has more than 200 publications in international peer-reviewed journals. She is the recipient of the DAE Science and Technology Excellence award 2017, DAE-SSPS Young Achievers Award (YAA) in 2008 and DAE Group Achievement award in 2008 for her work on gas sensors, and in 2014, for the growth of single crystals of CsI and NaI for radiation detection applications.

## Shashwati Sen

Scientific Officer/G Technical Physics Division, Physics Group Bhabha Atomic Research Centre (BARC), Trombay - 400 085, INDIA.

mong the conventional scintillators, CsI:TI is widely used for gamma spectrometry. Other than having a very high light yield, CsI:TI has an important ability of differentiating between the incoming radiation based on pulse shape discrimination (PSD). This differentiation arises from the presence of "short" and "long" decay component in the scintillation pulse. Divalent cation co-doping in CsI:TI are one of the ways of tailoring the electronic structure so as to increase its PSD ability and simultaneously reducing the afterglow.

In a recent study (Sisodiya et al, Optimizing the Scintillation Kinetics of Csl Scintillator Single Crystal by Divalent Cation Doping: Insights from Electronic Structure Analysis and Luminescence Studies, J. Phys. Chem. C, doi.org/10.1021/acs.jpcc.3c06098), experimental investigations along with electronic structure calculations were carried out to understand scintillation kinetics of pure Csl, and doped/co-doped with Tl, Sr and Eu. DFT based electronic structure calculations showed that DX like defect center are formed in the Csl:Tl matrix when co-doped with Sr<sup>2+</sup> and Eu<sup>2+</sup> which effects its scintillation kinetics. The study gave an understanding of how the defects need to be tailored to enhance the properties of Csl:Tl.